

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶ :

C07C 315/00

A2

(11) International Publication Number:

WO 99/47497

(43) International Publication Date: 23 September 1999 (23.09.99)

(21) International Application Number: PCT/CA99/00212

(22) International Filing Date: 12 March 1999 (12.03.99)

(30) Priority Data:

60/077.990	13 March 1998 (13.03.98)	US
9815856.1	21 July 1998 (21.07.98)	GB

(71) Applicant (for all designated States except US): MERCK FROSST CANADA & CO. [CA/CA]; 16711 Trans-Canada Highway, Kirkland, Quebec H9H 3L1 (CA).

(72) Inventors; and


(75) Inventors/Applicants (for US only): GAREAU, Yves [CA/CA]; (CA). LABELLE, Marc [CA/CA]; (CA). JUTEAU, Helene [CA/CA]; (CA). GALLANT, Michel [CA/CA]; (CA). LACHANCE, Nicolas [CA/CA]; (CA). BELLEY, Michel [CA/CA]; 16711 Trans-Canada Highway, Kirkland, Quebec H9H 3L1 (CA).

(74) Agent: MURPHY, Kevin, P.; Swabey Ogilvy Renault, Suite 1600, 1981 McGill College, Montreal, Quebec H3A 2Y3 (CA).

(81) Designated States: AE, AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published*Without international search report and to be republished upon receipt of that report.*

(54) Title: CARBOXYLIC ACIDS AND ACYLSULFONAMIDES, COMPOSITIONS CONTAINING SUCH COMPOUNDS AND METHODS OF TREATMENT



(57) Abstract

Compounds of formula (I), as well as pharmaceutically acceptable salts, hydrates and esters thereof, are disclosed. The compounds are useful for treating or preventing prostaglandin mediated diseases. Pharmaceutical compositions containing such compounds and methods of treatment are also included.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

5

CARBOXYLIC ACIDS AND ACYLSULFONAMIDES, COMPOSITIONS CONTAINING SUCH COMPOUNDS AND METHODS OF TREATMENT

BACKGROUND OF THE INVENTION

10

The present invention relates to compounds which are useful for treating or preventing prostaglandin mediated diseases, methods of treatment and pharmaceutical compositions containing such compounds. The compounds are structurally different from conventional NSAIDs and opiates, and are antagonists of the pain and inflammatory effects of E-type prostaglandins.

15

Two review articles describe the characterization and therapeutic relevance of the prostanoid receptors as well as the most commonly used selective agonists and antagonists: *Eicosanoids: From Biotechnology to Therapeutic Applications*, Folco, Samuelsson, Macclouf, and Velo eds, Plenum Press, New York, 1996, chap. 14, 137-154 and *Journal of Lipid Mediators and Cell Signalling*, 1996, 14, 83-87. An article from *The British Journal of Pharmacology* (1994, 112, 735-740) suggests that Prostaglandin E₂ (PGE₂) exerts allodynia through the EP₁ receptor subtype and hyperalgesia through EP₂ and EP₃ receptors in the mouse spinal cord.

20

25

Thus, selective prostaglandin ligands, agonists or antagonists, depending on which prostaglandin E receptor subtype is being considered, have anti-inflammatory, antipyretic and analgesic properties, and in addition inhibit hormone-induced uterine contractions. Moreover, the compounds have anti-cancer effects.


30

The compounds have a reduced potential for gastrointestinal toxicity, a reduced potential for renal side effects, a reduced effect on bleeding times and a lessened ability to induce asthma attacks in aspirin-sensitive asthmatic subjects.

35

5 SUMMARY OF THE INVENTION

The present invention relates to compounds represented by formula I:



I

10 as well as pharmaceutically acceptable salts, hydrates and esters thereof, wherein:

HET represents a 5-12 membered monocyclic or bicyclic aromatic ring system containing 0-3 heteroatoms selected from O, S(O)_n and N(O)_m wherein m is 0 or 1 and n is 0, 1 or 2;

15 A is a one or two atom moiety and is selected from the group consisting of: -W-, -C(O)-, -C(R⁷)₂-W-, -W-C(R⁷)₂-, -CR⁷(OR²⁰)-, -C(R⁷)₂-, -C(R⁷)₂-C(OR²⁰)R⁷-, -C(R⁷)₂-C(R⁷)₂- or -CR⁷=CR⁷-, wherein W represents O, S(O)_n or NR¹⁷, with n as previously defined and R¹⁷ as defined below;

20 X represents a 5-10 membered monocyclic or bicyclic aryl or heteroaryl group having 1-3 heteroatoms selected from O, S(O)_n and N(O)_m, and optionally substituted with R¹⁴ and R¹⁵, and A and B are attached to the aryl or heteroaryl group ortho relative to each other;

Y represents O, S(O)_n, NR¹⁷, a bond or -CR¹⁸=CR¹⁸-;

25 B represents - (C(R¹⁸)₂)_p-Y- (C(R¹⁸)₂)_q-

wherein p and q are independently 0-3, such that when Y represents O, S(O)_n, NR¹⁷ or -CR¹⁸=CR¹⁸-, p + q = 0-6, and when Y represents a bond, p + q is 1-6;

Z is OH or NHSO₂R¹⁹;

30 R¹ R² and R³ independently represent H, halogen, lower alkyl, lower alkenyl, lower alkynyl, lower alkenyl-HET(R^a)₄₋₉-, (C(R⁴)₂)_pSR⁵-, (C(R⁴)₂)_pOR³-, (C(R⁴)₂)_pN(R⁶)₂-, CN, NO₂-, (C(R⁴)₂)_pC(R⁷)₃-, CO₂R⁹-, CON(R⁶)₂ or - (C(R⁴)₂)_pS(O)_nR¹⁰-, wherein n and p are as previously defined;

35 each R⁴ is independently H, F, CF₃ or lower alkyl,

5 or two R^4 groups are taken in conjunction and represent a ring of up to six atoms, optionally containing one heteroatom selected from O, $S(O)_n$ or $N(O)_m$;

each R^5 is independently lower alkyl, lower alkenyl, lower alkynyl, CF_3 , lower alkyl-HET, lower alkenyl-HET or $-(C(R^{18})_2)_pPh(R^{11})_0-$
 10 2;

each R^6 is independently H, lower alkyl, lower alkenyl, lower alkynyl, CF_3 , Ph, Bn and when two R^6 groups are attached to N they may be taken in conjunction and represents a ring of up to 6 atoms, optionally containing an additional heteroatom selected from O, $S(O)_n$ or
 15 $N(O)_m$;

each R^7 is independently H, F, CF_3 or lower alkyl, and when two R^7 groups are presents, they may be taken in conjunction and represent an aromatic or aliphatic ring of 3 to 6 members containing from 0-2 heteroatoms selected from O, $S(O)_n$ and $N(O)_m$;

20 each R^8 represents H or R^5 ;

each R^9 is independently H, lower alkyl, lower alkenyl, lower alkynyl, Ph or Bn;

each R^{10} is independently lower alkyl, lower alkenyl, lower alkynyl, CF_3 , $Ph(R^{11})_0-3$, $CH_2Ph(R^{11})_0-3$ or $N(R^6)_2$;

25 each R^{11} is independently lower alkyl, SR^{20} , OR^{20} , $N(R^6)_2$, $-CO_2R^{12}$, $-CON(R^6)_2$, $-C(O)R^{12}$, CN, CF_3 , NO_2 or halogen;

each R^{12} is independently H, lower alkyl or benzyl;

each R^{13} is independently H, halo, lower alkyl, O-lower alkenyl, S-lower alkyl, $N(R^6)_2$, CO_2R^{12} , CN, CF_3 or NO_2 ;

30 R^{14} and R^{15} are independently lower alkyl, halogen, CF_3 , OR^{16} , $S(O)_nR^{16}$ or $C(R^{16})_2OR^{17}$;

each R^{16} is independently H, lower alkyl, lower alkenyl, Ph, Bn or CF_3 ;

each R^{17} is independently H, lower alkyl or Bn;

35 each R^{18} is independently H, F or lower alkyl, and when two R^{18} groups are present, they may be taken in conjunction and represent a ring of 3 to 6 members comprising carbon atoms and optionally one heteroatom chosen from O, $S(O)_n$ or N;

5 each R^{19} is lower alkyl, lower alkenyl, lower alkynyl, CF_3 ,
HET(R^a)₄₋₉, lower alkyl-HET(R^a)₄₋₉ or lower alkenyl-HET(R^a)₄₋₉;

 each R^{20} is independently H, lower alkyl, lower alkenyl,
lower alkynyl, CF_3 or $Ph(R^{13})_2$
and

10 each R^a is independently selected from the group consisting
of:

H, OH, halo, CN, NO₂, amino, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl,
C₁₋₆ alkoxy, C₂₋₆alkenyloxy, C₂₋₆alkynyloxy, C₁₋₆alkylamino,
di-C₁₋₆alkylamino, CF_3 , C(O)C₁₋₆alkyl, C(O)C₂₋₆alkenyl, C(O) C₂₋
15 6alkynyl, CO₂H, CO₂C₁₋₆alkyl, CO₂C₂₋₆alkenyl, and CO₂C₂₋₆alkynyl.

 said alkyl, alkenyl, alkynyl and the alkyl portions of
alkylamino and dialkylamino being optionally substituted with 1-3 of:
hydroxy, halo, aryl, C₁₋₆ alkoxy, C₂₋₆alkenyloxy, C₂₋₆alkynyloxy, CF_3 ,
C(O)C₁₋₆alkyl, C(O)C₂₋₆alkenyl, C(O)C₂₋₆alkynyl, CO₂H, CO₂C₁₋₆alkyl,
20 CO₂C₂₋₆alkenyl, CO₂C₂₋₆alkynyl, NH₂, NHC₁₋₆alkyl and N(C₁₋₆alkyl)₂.

Pharmaceutical compositions are also included which are
comprised of a compound of formula I in combination with a
pharmaceutically acceptable carrier.

 A method of treating or preventing a prostaglandin
25 mediated disease is also included which is comprised of administering
to a mammalian patient in need thereof, a compound of formula I in an
amount which is effective for treating or preventing a prostaglandin
mediated disease.

30 DETAILED DESCRIPTION OF THE INVENTION

 The present invention relates to carboxylic acids and
acylsulfonamides, which are ligands at prostaglandin receptors, as well
as a method for treating or preventing a prostaglandin mediated disease
comprising administering to a patient in need of such a treatment of an
35 amount of compound of Formula I which is effective for treating or
preventing a prostaglandin mediated disease.

 The invention described in this patent application is
described using the following definitions unless otherwise indicated.

5 HET represents a 5-12 membered aromatic ring system containing 0-3 heteroatoms selected from O, S(O)_n and N wherein n is 0, 1 or 2. HET may be substituted with up to three substituents on the aromatic ring system, R¹, R² and R³. "Aromatic ring systems" as used herein includes aryl and heteroaryl groups such as benzene,
10 naphthalene, biphenyl, pyridine, quinoline, isoquinoline, furan, benzofuran, thiophene, benzothiophene, oxazole, thiazole, imidazole, benzothiazole, triazole, 1,2,5-thiadiazole, thienopyridine, indole, tetrazole, imidazole, benzoxazole, 1,2-methylenedioxybenzene and pyrrole.

15 HET² is a subset of HET and represents a member selected from the group consisting of: phenyl, thienyl, naphthyl, furanyl, thiazolyl, imidazolyl and indolyl.

Aryl refers to aromatic 6-10 membered groups having 1-2 rings and alternating (resonating) double bonds. Examples include
20 phenyl, biphenyl and naphthyl.

Heteroaryl refers to aromatic 5-12 membered groups having alternating (resonating) double bonds and containing from 1-4 heteroatoms selected from O, S(O)_n and N. Examples include the following: : quinoline, furan, benzofuran, thiophene, benzothiophene,
25 thiazole, benzothiazole, 1,2,5-thiadiazole, thienopyridine, oxazole, indole, isoindole, pyridine, isoquinoline, imidazole, thiazole, triazole, 1,3-methylene dioxobenzene, pyrrole and naphthyridine,

Heterocyclyl refers to non-aromatic 5-12 membered cyclic groups having 1-4 heteroatoms selected from O, S(O)_n and N. Examples
30 of heterocyclic groups are piperidine, piperazine, pyrrolidine, tetrahydrofuran, tetrahydropyran and morpholine.

X represents a 5-10 membered monocyclic or bicyclic aryl or heteroaryl group having 1-3 heteroatoms selected from O, S(O)_n and N(O)_m, and optionally substituted with R¹⁴ and R¹⁵, and A and B are
35 attached to the aryl or heteroaryl group X in positions which are ortho relative to each other. Examples are selected from the group consisting of: phenyl, naphthyl, biphenyl, quinoline, furan, benzofuran, pyridyl, pyrrole, thiophene, benzothiophene, thiazole, benzothiazole, 1,2,5-

5 thiadiazole, triazole, 1,2-methylenedioxybenzene, thienopyridine, oxazole and indole.

The terms alkyl, alkenyl, and alkynyl mean linear, branched, and cyclic structures and combinations thereof.

"Lower alkyl" means alkyl groups of from 1 to 7 carbon
 10 atoms. Examples of lower alkyl groups include methyl, ethyl, propyl, cyclopropyl, isopropyl, butyl, s- and t-butyl, pentyl, cyclopentyl, hexyl, cyclohexyl, heptyl, and the like. When propyl and butyl are recited without the isomeric form being specified, these include all isomers thereof:

15 "Lower alkenyl" means alkenyl groups of 2 to 7 carbon atoms. Examples of lower alkenyl groups include vinyl, allyl, isopropenyl, pentenyl, hexenyl, heptenyl, 1-propenyl, 2-butenyl, 2-methyl-2-butenyl, cyclopropen-1-yl, cyclohexen-3-yl and the like. When
 20 cis or trans is not specified, both are intended in pure form as well as in the form of a mixture of isomers.

"Lower alkynyl" means alkynyl groups of 2 to 7 carbon atoms. Examples of lower alkynyl groups include ethynyl, propargyl, 3-methyl-1-pentynyl, 2-heptynyl, 2-(cyclopropyl)ethenyl, 3-(cyclobutyl)-1-propynyl and the like.

25 Halogen (halo) includes F, Cl, Br and I.

The following abbreviations have the indicated meanings:

	AIBN	=	2,2'-azobisisobutyronitrile
	B.P.	=	benzoyl peroxide
	Bn	=	benzyl
30	CCl ₄	=	carbon tetrachloride
	D	=	-O(CH ₂) ₃ O-
	DAST	=	diethylamine sulfur trifluoride
	DCC	=	dicyclohexyl carbodiimide
	DCI	=	1-(3-dimethylaminopropyl)-3-ethyl
35			carbodiimide
	DEAD	=	diethyl azodicarboxylate
	DIBAL	=	diisobutyl aluminum hydride
	DME	=	ethylene glycol dimethylether
	DMAP	=	4-(dimethylamino)pyridine
40	DMF	=	N,N-dimethylformamide
	DMSO	=	dimethyl sulfoxide
	Et ₃ N	=	triethylamine
	LDA	=	lithium diisopropylamide

5 HET represents a 5-12 membered monocyclic or bicyclic aromatic ring system containing 0-3 heteroatoms selected from O, S(O)_n and N(O)_m wherein m is 0 or 1 and n is 0, 1 or 2;

A is a one or two atom moiety and is selected from the group consisting of: -W-, -C(O)-, -C(R⁷)₂-W-, -W-C(R⁷)₂-, -CR⁷(OR²⁰)-,
 10 -C(R⁷)₂-, -C(R⁷)₂-C(OR²⁰)R⁷-, -C(R⁷)₂-C(R⁷)₂ or CR⁷=CR⁷, wherein W represents O, S(O)_n or NR¹⁷, with n as previously defined and R¹⁷ as defined below;

X represents a 5-10 membered monocyclic or bicyclic aryl or heteroaryl group having 1-3 heteroatoms selected from O, S(O)_n and
 15 N(O)_m, and optionally substituted with R¹⁴ and R¹⁵, and A and B are attached to the aryl or heteroaryl group ortho relative to each other;

Y represents O, S(O)_n, NR¹⁷, a bond or -CR¹⁸ = CR¹⁸-;

B represents - (C(R¹⁸)₂)_p-Y- (C(R¹⁸)₂)_q-

wherein p and q are independently 0-3, such that when Y represents O,
 20 S(O)_n, NR¹⁷ or -CR¹⁸ = CR¹⁸-, p + q = 0-6, and when Y represents a bond, p + q is 1-6;

Z is OH or NHSO₂R¹⁹;

R¹ R² and R³ independently represent H, halogen, lower alkyl, lower alkenyl, lower alkynyl, lower alkenyl-HET(R^a)₄₋₉, -
 25 (C(R⁴)₂)_pSR⁵, -(C(R⁴)₂)_pOR⁸, -(C(R⁴)₂)_pN(R⁶)₂, CN, NO₂, -(C(R⁴)₂)_pC(R⁷)₃, -CO₂R⁹, -CON(R⁶)₂ or -(C(R⁴)₂)_pS(O)_nR¹⁰, wherein n and p are as previously defined;

each R⁴ is independently H, F, CF₃ or lower alkyl,

or two R⁴ groups are taken in conjunction and represent a ring of up to
 30 six atoms, optionally containing one heteroatom selected from O, S(O)_n or N(O)_m;

each R⁵ is independently lower alkyl, lower alkenyl, lower alkynyl, CF₃, lower alkyl-HET, lower alkenyl-HET or -(C(R¹⁸)₂)_pPh(R¹¹)₀₋₂.

35 each R⁶ is independently H, lower alkyl, lower alkenyl, lower alkynyl, CF₃, Ph, Bn and when two R⁶ groups are attached to N they may be taken in conjunction and represents a ring of up to 6 atoms,

R ¹ R ² R ³ -Het	A	X	B	Cpd
2-(benzo[b]thiophenyl)	CH ₂	4-F-1,2-Ph	CH=CH	539
5-(1-benzyl)indolyl	CH ₂	4-F-1,2-Ph	CH=CH	540
1-(6-(4-chloro)phenyl) indolyl	CH ₂	4-F-1,2-Ph	CH=CH	541
1-(5-chloro)indolyl	CH ₂	3,2-Pyr	CH=CH	542

5

wherein D = -O(CH₂)₃-O, Qn = 7-chloroquinolin-2-yl, 1,2-Ph = 1,2-benzenediyl, Rs = -CH₂SCH₂CH₂Ph, Pyr = pyridinediyl, c-pr = cyclopropyl and Bn = benzyl.

19. A pharmaceutical composition which is
 10 comprised of a compound in accordance with any one of claims 1 to 18 in combination with a pharmaceutically acceptable carrier.

20. A method of treating or preventing a prostaglandin mediated disease which is comprised of administering to a mammalian
 15 patient in need of such treatment a compound in accordance with claim 1 in an amount which is effective for treating or preventing a prostaglandin mediated disease.

21. A method in accordance with claim 19 wherein the
 20 prostaglandin mediated disease is selected from the group consisting of:
 pain, fever or inflammation associated with rheumatic fever, influenza or other viral infections, common cold, low back and neck pain, skeletal pain, post-partum pain, dysmenorrhea, headache, migraine, toothache, sprains and strains, myositis, neuralgia,
 25 synovitis, arthritis, including rheumatoid arthritis, degenerative joint diseases (osteoarthritis), gout and ankylosing spondylitis, bursitis, burns including radiation and corrosive chemical injuries, sunburns, pain following surgical and dental procedures, immune and autoimmune diseases;
 30 cellular neoplastic transformations or metastatic tumor growth;
 diabetic retinopathy, tumor angiogenesis;

- 5 prostanoid-induced smooth muscle contraction associated
with dysmenorrhea, premature labor, asthma or eosinophil related
disorders;
- Alzheimer's disease;
 glaucoma;
10 bone loss;
 osteoporosis;
 promotion of bone formation;
 Paget's disease;
 cytoprotection in peptic ulcers, gastritis, regional enteritis,
15 ulcerative colitis, diverticulitis or other gastrointestinal lesions; GI
bleeding and patients undergoing chemotherapy;
 coagulation disorders selected from hypoprothrombinemia,
haemophilia and other bleeding problems;
 kidney disease;
20 thrombosis;
 occlusive vascular disease;
 presurgery;
 and anti-coagulation.
- 25 22. A method in accordance with claim 20 wherein the
prostaglandin mediated disease is selected from the group consisting of:
pain, fever or inflammation.
23. A method in accordance with claim 20 wherein the
30 prostaglandin mediated disease is dysmenorrhea.
24. A method in accordance with claim 20, wherein the
compound is co-administered with other agents or ingredients.
- 35 25. A method in accordance with claim 24 wherein the
compound I is co-administered with another agent or ingredient
selected from the group consisting of: an analgesic selected from
acetaminophen, phenacetin, aspirin, a narcotic;

- 5 a COX-2 selective NSAID and a conventional NSAID;
caffeine;
an H₂-antagonist;
aluminum or magnesium hydroxide;
simethicone;
10 a decongestant selected from phenylephrine,
phenylpropanolamine, pseudophedrine, oxymetazoline, ephinephrine,
naphazoline, xylometazoline, propylhexedrine, or levo-desoxyephedrine;
an antiitussive selected from codeine, hydrocodone,
caramiphen, carbetapentane and dextramethorphan;
15 another prostaglandin ligand selected from misoprostol,
enprostil, rioprostil, ornoprostol and rosaprostol; a diuretic; and
a sedating or non-sedating antihistamine.

26. Use of a compound, salt, hydrate or ester as
defined in any one of claims 1 to 18 in the manufacture of a
20 medicament for treatment or prevention of a prostaglandin
mediated disease.

27. A compound, salt, hydrate or ester as defined in
any one of claims 1 to 18 for use in the treatment or prevention of
a prostaglandin mediated disease.

25 28. A prostaglandin antagonist pharmaceutical
composition comprising an acceptable prostaglandin antagonistic
amount of a compound, salt, hydrate or ester as defined in any one
of claims 1 to 18, in association with a pharmaceutically
acceptable carrier.